

## **PERFORMANCE AND ANALYSING CHARACTERISTICS IN DIFFERENT AGE GROUP OF KHO KHO PLAYERS**

NunavathRajitha

rajithanunavath123@gmail.com

Physical Director, Alphoras Degree PG College,  
Karimnagar ,Telangana.

### **ABSTRACT:**

In India, khokho is one of the oldest and most well-known tag games. The purpose of the study was to compare the physical fitness parameters and anthropometric parameters among the three age groups of Kho-Kho players as physical fitness and anthropometric parameters are the most important factors to execute performance in competitive situation. To accomplish the purpose of the study BMI, Arm length, Foot length, speed, agility and speed endurance were taken as the variables. To conduct the study total 45 girls subjects were selected from three different age group of Kho-Kho. The mean and S.D. of scores of each variables were computed, thereafter „F“ test was computed to find out the significance of differences among the score of each variable of physical fitness and anthropometric parameters among three groups of Kho-Kho players and LSD was computed to find out the significant difference among three groups. There was a gradual increase of height and body weight with age. The BMI of the junior group( $18.03 \pm 2.55 \text{ kg/m}^2$ ) was the lowest and the most senior group( $20.02 \pm 1.57 \text{ kg/m}^2$ ) was highest. The speed of the middle age group( $7.69 \pm 0.34\text{sec}$ ) was best. The agility of the junior group( $10.97 \pm 3.07\text{sec.}$ ) showed best than the other groups. The speed endurance of middle group( $81.13 \pm 7.15\text{sec}$ ) was better than the other two groups. The average foot length of junior most groups ( $21.86 \pm 1.24\text{cm}$ ) was lowest and the middle aged group( $22.86 \pm 1.12\text{cm}$ ) was highest. The average arm length of junior most group( $62.80 \pm 4.17\text{cm}$ ) lowest and the senior most group( $69.53 \pm 4.85\text{cm}$ ) was highest. It may conclude that the characteristics of anthropometric parameters and physical fitness variables vary with the different age but no such specific direction of increase or decrease of such variables is drawn.



## INTRODUCTION

A highly regarded indigenous sport in India is kho-kho. [23][15][6] In India, khokho is the most widely practised traditional sport.Kho-Kho is a very intricate and strategic sport.Kho-Kho is a tag sport in which teams of twelve players attempt to avoid being touched by members of the opposing team; only nine players from each team enter the field. It is one of the two traditional tag games most played in South Asia, along with Kabaddi. It is played elsewhere besides South Asia (primarily Bangladesh, India, and Pakistan).[22] KhoKho's ancestry is difficult to determine, but many historians think that it is a modified version of the game "Run Chase," which in its most basic form entails chasing and touching.[12] Fast and furious action is needed in khokho.[8] Physical fitness is the state of having the optimum possible functioning of the heart, blood vessels, lungs, and muscles.[9] The primary goal of sports training is the enhancement and maintenance of physical fitness.[23] Physical fitness forwards an athletes for better athletic performance. Athletes should develop the ability to play the game with good skill and perform consistently well.[24] Naturally kho-kho game is vigorous and combative types. For excellent execution of the kho-kho skills like dodging, feinting, and bursts of speed players at individual level requires the variables of stamina, endurance, strength, and agility.[12] For develop excellent physical Fitness, Kho-Kho players requires important variables like Speed, Explosive Strength, VO<sub>2</sub> Max, Agility, Flexibility etc.[10] In quick games and sports like basketball and khokho, agility and speed are crucial elements. Players on both the offensive and defensive sides can change directions quickly and with good speed, but switching positions is

uncommon.[18] Due to the need of swift movements in confined spaces, kho-kho games require excellent agility, explosive strength, and endurance.[28] Age is a crucial biomarker that reveals athletes' performance traits. Physical function changes as people age. Different games may be appropriate for different ages.[2] Performance is determined by endogenous variables like anthropometric, genetic and physiological traits.[16][21] Several researchers hypothesised that various body types, shapes and proportions are advantageous for various types of physical activity.[17][13][24] A number of anthropometric traits, including body composition and somatotype, have been linked to athletic success.[7][11] An athlete's ideal physique, successful participation and level of performance for a certain sport may be determined by their anthropometric and physical attributes.[14] "Body types" play an important role in sports. Anthropometric measurements have shown a relationship between physical attributes, body type, and athletic ability. Height, weight, and other anthropometric factors are extremely important to player performance in all games. The most crucial deciding factors in winning games are anthropometric characteristics and physical prowess, particularly when both teams' skill and planning levels are comparable. Anthropometric measurements are important for determining weight and height at various ages and for assessing one's growth.[5] In many games, physical structure, particularly height and arm length, provides a clear and decisive advantage. Similarly, segmental length of individual body parts, specifically arm length, provides a significant advantage in certain athletic events and games.[29] The ability to balance, move quickly, and direct voluntary movements of people toward a goal are all physical

traits that are related to the feet, particularly when changing directions and maintaining balance while using attacking and defending techniques. When performing fundamental motions like standing, running, and jumping, the length and width of the foot are very important.[5] Some specific anthropometric traits are necessary to succeed in a sporting event. There are certain distinctions between athletes' physical structures and composition in respective to team and individual sports.[19] The ability to perform daily tasks may be limited by an individual's physical and physiological characteristics, including skill performance and anthropometric variables. But in sports field efficiency of the execution of movement and sustainable capacity may not be similar for athletes with same height and weight. Consequently, segments of the body part vary in terms of their length, breadth, and circumference.[31] Anthropometric and Physical performance measurements of running, hop-testing, and jumping were found to have a significant and relevant association.[1] There are appropriate research-based criteria for selecting standard players, which are also lacking in KhoKho. Research-based experience is always beneficial in the development of various games and sports, which is also important for 'KhoKho' field.[8] The aim of present study is to find out the selected anthropometric determinants and physical fitness variable as the performance Characteristics of Kho-Kho and to compare them among three different age groups.

## METHODOLOGY

**Materials and Methods:- Selection of Subjects:** To conduct the study total 45 girls Kho-Kho players were taken and their age ranges between 14 to 23 years. All the subjects were selected from BandelKodialiaSangha, KheyaliSangha, Hooghly and Jadavpur University. All

subjects were equally divided into three groups. Each group had possessed with 15 players and the groups were namely Gr-A, Gr-B and Gr-C. The Gr-A was of 14-16 years, Gr-B of 16-18 years and Gr-C of 18-23years. Selections of variables:

B.M.I, Foot length and Arm length were taken as Anthropometric variables and Speed, Agility and Speed Endurance were taken as Physical Fitness variables. Design of the study: All the anthropometric measurements were taken such as height and weight for BMI, Arm length and foot length etc. After that the test of 50 meter dash and 4×10 mtrshuttle run were taken to measure speed and agility as fitness variables. Then 30 minutes rest has been given. After 30 minutes rest 400 meter run were conducted to measure speed endurance. Statistical Technique: The mean and S.D. of scores of each variables were computed, there after „F“ test was computed to find out the difference among the score of each variable of physical fitness and anthropometric parameters among three groups of Kho-Kho players, the LSD was computed to find out the significant difference among three groups. The level of significance was set at 0.05 level ( $p<0.05$ ).

## RESULTS:

Table no. 1:- Mean, S.D. of age, height, weight and BMI of different Age Groups.

Groups	Mean±S.D. of age (yrs.)	Mean±S.D. of height (cm.)	Mean±S.D. of weight (kg.)	Mean±S.D. BMI(kg/mtr. <sup>2</sup> )
A	15.20 ± 0.41	144.73 ± 8.97	38.06 ± 7.94	18.03 ± 2.55
B	17.33 ± 0.48	153.06 ± 8.31	44.53 ± 5.04	19.04 ± 2.07
C	21.20 ± 1.97	153.46 ± 6.03	46.60 ± 3.68	20.02 ± 1.57

It was observed from table-1, that the Mean & S.D. of age of group A was  $15.20 \pm 0.41$  yrs., group B was  $17.33 \pm 0.48$  yrs. and group C was  $21.20 \pm 1.97$  yrs. From the above table it was clear that the Mean and S.D. of height of Group A was  $144.73 \pm 8.97$  cm., group B was  $153.06 \pm 8.31$  cm. and group C was  $153.46 \pm 6.03$  cm. It has

observed that there was a linear progression of height with age. This was very inevitable as the height increases with the anatomical growth and that continued generally up to the age of 18yrs+ & above in case of girls. It has indicated that the Mean and S.D. of weight of group A was  $38.06 \pm 7.94\text{kg.}$ , group B was  $44.53 \pm 5.04\text{kg.}$  and group C was  $46.60 \pm 3.68\text{kg.}$  There was a gradual increase of body weight also with age. It was simply due to the growth pattern of the considered age group. The Mean and S.D. of BMI of the group A was  $18.03 \pm 2.55\text{ kg./mt.2}$ , the group B was  $19.04 \pm 2.07\text{ kg./mt.2}$  and the group C was  $20.02 \pm 1.57\text{ kg./mt.2}$ . The Mean score of the BMI of group A was lowest and the group C was highest.

**Table no. 2:-** ANOVA on BMI Component of Different Age Groups

Source	Sum square	d. f.	Mean square variance	'p' value	F-Ratio
Between Gr.	29.663	2	14.831		
Within gr.	185.969	42	4.428	0.045*	3.350
Total	215.632	44			

Level of significance at 0.05%

Table-2 shows that there was significant difference existed among the groups as the „p“ value was less than the significance level 0.05. Hence to found the relative comparativeness between the said 3 groups and to understand the difference between groups the LSD Post Hoc test were applied.

**Table no. 3:-** Post Hoc Tests (LSD), Multiple Comparisons of BMI.

Group (I)	Group (J)	Mean Difference (mean of group I- Mean of group J)	Std. Error	Sig.	95% confidence Interval		
					Lower Bound	Upper Bound	
14'-16 Yrs.	A	B	-1.00733	0.76836	0.197	-2.5579	0.5433
	C		-1.98867	0.76836	0.013**	-3.5393	-0.4381
16'-18 Yrs.	B	C	-0.98133	0.76836	0.209	-2.5319	0.5693

BMI is an energy indicator that correlates total mass and height, allowing athletes to be compared across varied distances.[27]It has seen from

table-3, that there was significant difference existed only between group A and C. It has reported similarly that the BMI increased significantly with the increase of age and this result almost support the present investigation. Body mass index above  $23\text{ kg/m}^2$  was strongly correlated with nearly all Kho-Kho abilities, with an overall coefficient of correlation was 0.83.[8] Marc et al study's shows an optimum BMI of  $19.8\text{ kg/m}^2$ , despite the fact that the top ten performers of all time have BMIs ranging from  $17.5$  to  $20.7\text{ kg/m}^2$ .

Table no. 4:- Mean, S.D. of foot length, arm length, speed, agility, speed endurance of different Age Groups.

Groups	Mean $\pm$ S.D. of foot length (cm.)	Mean $\pm$ S.D. of arm length (cm.)	Mean $\pm$ S.D. of speed (sec.)	Mean $\pm$ S.D. of agility(sec.)	Mean $\pm$ S.D. of speed endurance(sec.)
A	$21.86 \pm 1.24$	$62.80 \pm 4.17$	$8.12 \pm 0.38$	$10.97 \pm 3.07$	$101.33 \pm 5.85$
B	$22.86 \pm 1.12$	$68.26 \pm 3.30$	$7.69 \pm 0.34$	$11.33 \pm 0.68$	$81.13 \pm 7.15$
C	$22.80 \pm 1.08$	$69.53 \pm 4.85$	$8.34 \pm 0.31$	$11.34 \pm 0.65$	$97.53 \pm 6.18$

The Mean and S.D. of foot length of the group A was  $21.86 \pm 1.24\text{ cm.}$ , the group B was  $22.86 \pm 1.12\text{ cm}$  and the group C was  $22.80 \pm 1.08\text{ cm.}$  The Mean score of the foot length of group A was lowest and the group B was highest. From table value it was shows that there was a difference in foot length among the groups. The significance of statistical differences in foot length among the groups was assessed by applying “F” test.

**Table no. 5:- ANOVA on Foot Length Component of Different Age Groups.**

Source	Sum square	d. f.	Mean square	'p' value	F-Ratio
Between Gr.	9.378	2	4.689		
Within gr.	55.867	42	1.330	0.038**	3.525
Total	65.244	44			

Level of significance at 0.05%

From table- 4; it was cleared that there was significant difference existed among the groups as the „p“ value was less than the significance level 0.05. Hence to found the relative comparativeness between the said 3 groups and to understand the difference between groups the LSD Post Hoc test were applied. Table no. 6:- Post Hoc Tests (LSD), Multiple Comparisons of Foot length.

Group (I)	Group (J)	Mean Difference (mean of group I - Mean of group J)	Std. Error	Sig.	95% confidence Interval	
					Lower Bound	Upper Bound
14'-16 Yrs.	B	-1.00000	0.42113	0.022**	-1.8499	-0.1501
	C	-0.93333	0.42113	0.032**	-1.7832	-0.0834
16'-18 Yrs.	B	0.06667	0.42113	0.875	-0.7832	0.9166
	C					

Significant differences were found between group A and B & between group A and C.Darwish in his study found that foot length, transitional speed test, and agility all significantly correlate with one another.Nearly all KhoKho skill exhibited a strong link with long leg length (40 inches and above), and the total coefficient of correlation was 0.82. The Mean and S.D. of arm length of the group A was  $62.80 \pm 4.17$ cm., the group B was  $68.26 \pm 3.30$ cm. and the group C was  $69.53 \pm 4.85$ cm. The Mean score of the arm length of group A was lowest and the group C was highest.Singh&Jaiswal, (2016)in their study also found almost similar results

about arm length i.e. $72.54 \pm 3.6$ cm (ages spanning 17 to 26 years).

**Table no. 7:- ANOVA on Arm Length Component of Different Age Groups.**

Source	Sum square	d. f.	Mean square	'p' value	F-Ratio
Between Gr.	384.133	2	192.067		
Within gr.	727.067	42	17.311	0.000**	11.095
Total	1111.200	44			

Level of significance at 0.05%

From table- 7; it was cleared that there was significant difference observed among the groups of Kho-Kho players at 0.05 levels in respect of arm length. To understand the difference between groups the LSD Post Hoc test were applied. Table no. 8:- Post Hoc Tests (LSD), Multiple Comparisons of Arm length

Group (I)	Group (J)	Mean Difference (mean of group I - Mean of group J)	Std. Error	Sig.	95% confidence Interval	
					Lower Bound	Upper Bound
14'-16 Yrs.	B	-5.46667	1.51926	0.001**	-8.5327	-2.4007
	C	-6.73333	1.51926	0.000**	-9.7993	-3.6673
16'-18 Yrs.	B	-1.26667	1.51926	0.409	-4.3327	1.7993
	C					

Significant at .05 level

It has seen that there was a significant difference existed between group A and B & group A and C but there was no significant difference between group B and C. KhoKho skills were shown to have overall coefficients of correlation of 0.75, 0.59, 0.68, 0.65, and 0.68 with regard to height, weight, BMI, arm length, and leg length. The findings showed a significant ( $p<.01$ ) relationship between all of the anthropometric factors and KhoKho abilities. It has been observed that having a long arm (32 inches or more) has a high link with almost all KhoKho skills, with an overall coefficient of



correlation of 0.80.[8] The mean and S.D. of speed of the group A was  $8.12 \pm 0.38$  sec., group B was  $7.69 \pm 0.34$  sec. and group C was  $8.34 \pm 0.31$  sec. The speed of the middle age group B was maximum. Table no. 9:- ANOVA on Speed Component of Different Age Groups

Source	Sum square	d. f.	Mean square variance	'p' value	F-Ratio
Between gr.	3.248	2	1.624	0.000*	13.371
Within gr.	5.101	42	0.121		
Total	8.348	44			

Level of significance at 0.05%

From table- 9; it was cleared that there was significant difference existed among the groups as the „p“ value was less than the significance level 0.05. Hence to found the relative comparativeness between the said 3 groups and to understand the difference between groups the LSD Post Hoc test were applied. Table no. 10:-Post Hoc Tests (LSD), Multiple Comparisons of speed.

Group (I)	Group (J)	Mean Difference (mean of groupJ-Mean of groupI)	Std. Error	Sig.	95% confidence Interval
14'-	Gr.A	0.42600	0.12725	0.002*	0.1692 - 0.6828
16 Yrs.	B	-0.22133	0.12725	0.089	-0.4781 - 0.0355
16'-18 Yrs.	Gr. B	-0.64733	0.12725	0.000*	-0.9041 - 0.3905

Significant at .05 level

It has seen from the above table that the mean difference between group A and B was significant and between group B and C was significant. In the table of ANOVA it has declared that there were significant differences in case of speed of these three groups. We know that speed is stimulated by leg explosive strength and the muscle attached with pelvis which indicates that the maturity of muscles growth is very important and it is mainly a genetic factor. The speed is

measured by stride length and stride frequency and the stride length is largely influenced by the lower limb length. At this age group, in the late adolescents" period the growth pattern took maximum spurt with sudden increase of strength. All these factors are very much boosting in case of speed performance. In terms of speed, Sahu D. P., (2019) observed that national level KhoKho player exhibit greater sprinting performance when compared to state level KhoKho players.[26] But the present researcher has not found any such speed related result when taking into account their age group. The Mean and S.D. of the time taken to cover  $4 \times 10$  mtr., shuttle run, which was indicated the agility of the group A was  $10.97 \pm 3.07$  sec. and group B was  $11.33 \pm 0.68$  sec. and group C was  $11.34 \pm 0.65$  sec. The mean score of the agility component of group A was better followed by group B and group C. Singh & Jaiswal, (2016), in their study also found almost similar results about agility i.e.  $12.1 \pm .65$  sec. (ages spanning 17 to 26 years & agility run test was of  $4 \times 15$  mtr.).[28] Table no.11:- ANOVA on agility component of different age groups

Source	Sum square	d. f.	Mean square variance	'p' value	F-Ratio
Between Gr.	1.315	2	0.658		
Within gr.	144.916	42	3.450	0.827	0.191
Total	146.231	44			

Level of significance at 0.05%



From the above table of ANOVA it has clear that there was no significant difference in case of agility among those three groups. We know that agility is a component which is a combination of speed and quick turning ability. These two factors again stimulated by leg explosive strength and the muscle of pelvis which indicate that the maturity of muscles growth is very important. Here the present researcher considered the three groups whose age totally ranged between 14 to almost 23 yrs. In this age group the anatomical structure almost became matured and the Physiological functioning became maximum. As the participants were from the village area so naturally they are habituated enough with high intensity movement as considering their daily life style. Hence considering the said different factors it is clear that, in said specific age range there were a little possibility to differ in case of a combined effect of speed in a very short distance and turning ability as also they are mainly stimulated by genetic factor. Higher agility players (7.96 seconds and lower) had strong relationships with almost all Kho-Kho skills, with an overall coefficient of correlation of 0.86.[8] In a different study in the area of agility Sahu D. P., (2019) observed that national level KhoKhoplayer exhibit

better agility performance in comparison to state level KhoKho players.[26] Female khokho players at different levels were shown to have significant differences in their agility, which gradually increases from the sub-junior group to the senior group, according to Roy et al. (2016).[23] The Mean and S.D. of the time taken to cover 400mtr., which indicates the speed endurance of the group A was  $101.33 \pm 5.85$  sec., group B was  $81.13 \pm 7.15$  sec. and group C was  $97.53 \pm 6.18$  sec. The Mean score of the speed endurance component of group B was better followed by group C and Group A. Table no.12:- ANOVA on Speed Endurance Component of Different Age Groups.

Source	Sum square	d. f.	Mean square variance	'p' value	F-Ratio
Between Gr.	3457.200	2	1728.600		
Within gr.	1730.800	42	41.210	0.000**	41.947
Total	5188.000	44			

Level of significance at 0.05%

It shows that there was significant difference observed among the three groups of Kho-Kho players at 0.05 levels. So, to understand the difference between groups the LSD Post Hoc test was applied. Table no. 13:- Post Hoc Tests (LSD), Multiple Comparisons of Speed endurance

Group ( I )	Group(J)	Mean Difference (mean of group I -	Std. Error	Sig.	95% confidence Interval	
					Lower	Upper
14'-16 Yrs	B	20.20000	2.34406	0.000**	15.4695	24.9205
	C	3.80000	2.34406	0.112	-0.9305	8.5305
16'-18 Yrs	C	-16.40000	2.34406	0.000**	-21.1305	-11.6695
	B					

Significant at .05 level

In the table of ANOVA it has declared that there were significant differences in



case of speed endurance of these three groups. We know that speed endurance is stimulated by leg explosive strength and through the aerobic capacity which indicate that the maturity of muscles growth is very important and it is mainly a genetic factor. At this age group, in the late adolescents" period the growth pattern took maximum spurt with sudden increase of strength. All these factors are very much boosting in case of speed endurance performance.Ghosh et.al (2014) in a different study found that the overall coefficients of correlation between KhoKhoskillswith cardio-respiratory endurance,agility and 50m dash were 0.58, 0.60, and 0.65 respectively and suggested that all the selected physical variables were significantly( $p < .01$ )related with the KhoKhoskills.

## CONCLUSION :

On the basis of result obtain it has concluded that there was a gradual increase of height and body weight with age.The speed of the middle age group was best than the other two groups.The agility of the junior most group showed better than their senior groups.The speed endurance of middle group was better than the other two groups. The BMI of the senior group was the highest and the

junior most group was lowest. The average foot length of junior most group was lowest and the middle aged group was highest.The average arm length of junior most group lowest and the senior most group was highest. Finally, the present study concludes that specific age groups are characterized with some specific anthropometric and fitness variables. This study provides reference values of anthropometric characteristics and physical fitness status about the different age group kho-kho players which may be informative for coaches to frame and control the training process in order to enhance talent identification in khokho, as well as players" performances.

## REFERENCES:

1. Baacke, L.W. (1984). Relationship of Selected Anthropometric and Physical Performance Measures to Performance in Running Hop-Step and Jump, *Research Quarterly*, 35: 1984 2.
- Berthelot G., Johnson S., Noirez P., Antero J., Marck A., Desgorges F. D., Pifferi F., Carter P. A., Spedding M., Manoux A. S. & Toussaint J. F. (2019). The age-performance relationship in the general population and strategies to delay age related decline in performance. *Archives of Public Health* volume 77, no. 51

- 3.Canhadas, Ive& Silva, Rodrigo &Chaves, Celso&Portes, Leslie.(2010). Anthropometric and physical fitness characteristics of young male soccer players.*Revista Brasileira de Cineantropometria e Desempenho Humano.* 12. 4. Damsgaard R, Bencke J, Matthiesen G, Petersen JH, Müller J. (2001). Body proportions, body composition and pubertal development of children in competitive sports.*Scand J Med Sci Sports;*11:54-60
5. Darwish, N. H. (2015). The relation between foot's length & width with some physical abilities and performance accuracy of some basic skills of female volleyball players.*The Swedish Journal of Scientific Research.*Vol. 2. Issue 5. 6. Dabholkar, A. (2017). Evaluation of foot posture and Correlation with the injury in female KhoKho players. *International Journal of Physical Education, Sports and Health,* 4, 381–384.
7. Eston, R. Reilly, T. (2001), *Kinanthropometry and exercise physiology laboratory manual: Tests, Procedure and data.*, London and New York: Routledge, Tayler and Francis Group. 2 ed., Vol. 1, pp. 47-65
- 8.Ghosh, M., &Kundu, B. (2014).Physical, Physiological and Anthropometric Measures as Determinants of Performance in Kho-Kho Skills - A Correlational Study.*International Journal of Humanities and Social Science Invention.* Volume 3 Issue 8, PP 04-12 9.
- Getchell, Bud, *Physical Fitness: A Way of Life*, New York: Willey and Sons, inc., 1979 10. Garai, Biswajit&Biswas, Sudarsan.(2017). A Comparative Profile Study of West Bengal University Level KhoKho Players. 4. 421-428. 11. Heath B. H. and Carter J.E.L. (1990). Somatotyping development and application.*Cambridge University press*, New York. 12.Jaiswal, A. (2014). Anthropometric and somatotyping study among the female Kho-Kho players of Pondicherry: A comparative analysis. *J. Glob. Econ.*, 2, 1–3. Scopus.

**NunavathRajitha :**

I NunavathRajitha ,presently working as Physical Director in Alphores Women's Degree and PG college, karimnagar.I have completed MBA in sathavahana University in 2012. I have completed BPED in Barkatullah University in 2014.I have completed MPED in Vikramasimhapuri University, Nellore with 73% in 2018 . I have participated in many games in schools and college levels and got many medals.